



## Maize Pests in Mexico and Challenges for the Adoption of Integrated Pest Management Programs

Carlos A. Blanco,<sup>1,2</sup> José; Guadalupe Pellegaud,<sup>3</sup> Urbano Nava-Camberos,<sup>4</sup> David Lugo-Barrera,<sup>5</sup> Paulina Vega-Aquino,<sup>3</sup> Jesús Coello,<sup>6</sup> Antonio P. Terán-Vargas,<sup>7</sup> AND Jesús Vargas-Camplis<sup>7</sup>

<sup>1</sup>Biology Department, The University of New Mexico, Albuquerque, NM 87131.

<sup>2</sup>Corresponding author, e-mail: cblancom@unm.edu.

<sup>3</sup>Ejido 18 de marzo, Los Mochis, Sinaloa, Mexico.

<sup>4</sup>Universidad Juárez del Estado de Durango, Av. Universidad s/n Gómez Palacio, Durango, Mexico.

<sup>5</sup>Dow AgroSciences Mexico, Zapopan, Jalisco, Mexico.

<sup>6</sup>Grupo MASECA, Ocozocoautla de Espinoza, Chiapas, Mexico.

<sup>7</sup>INIFAP Tamaulipas, Cuauhtémoc, Tamaulipas, Mexico.

J. Integ. Pest Mngmt. 5(4): 2014; DOI: <http://dx.doi.org/10.1603/IPM14006>

**ABSTRACT.** Mexico is the fifth largest maize producer and the area of origin and domestication of this crop. Mexico's 6.8 million hectare annual production of maize represents 30% of its total domestic agricultural production and 6.6% of the country's arable land. However, Mexico's average yield of 3.17 tons per hectare is 38% below the world's average. Although maize is a major crop globally, it is typically not grown under modern or comprehensive integrated pest management (IPM) programs aimed at minimizing economic damage and lowering environmental and health risks. Pest management in commercial maize in Mexico continues to rely primarily on chemical control using conventional synthetic insecticides. Approximately 3,000 tons of active ingredient per year are used to combat the single most important pest, the fall armyworm (*Spodoptera frugiperda* Smith), followed by the corn earworm (*Helicoverpa zea* Boddie) and the black cutworm (*Agrotis ipsilon* (Hufnagel)). Mexican farmers have successfully adopted IPM programs for many crops, but IPM is not common in maize crops. Although different components and elements have been identified, most of the information on maize pests and IPM in Mexico has been published in technical bulletins of a reduced regional scope or in difficult-to-access reports, generally describing laboratory screenings, preliminary field evaluations, or listing technical recommendations but rarely including products and practices that growers actually use. This report provides up-to-date information on the common pests in the main maize-growing areas of Mexico from experts in the field. Updated information on maize pest population dynamics and control has been gathered from agricultural professionals directly involved in maize crop protection. This critical information, not readily available for Mexican production systems, is intended to help researchers, government officials, and industry specialists in making decisions regarding time and resource allocations in the design and implementation of IPM techniques, practices, and programs for Mexican maize.

**RESUMEN.** México es el centro de origen y domesticación del maíz y el quinto mayor productor de este grano del mundo, ocupando más de 6% de su superficie arable, lo que representa una tercera parte de su producción agrícola interna. El manejo integrado de plagas (MIP) puede ser una estrategia importante para reducir el daño de plagas y disminuir la cantidad de insecticida que se usa. El problema de plagas en este cultivo lo encabeza *Spodoptera frugiperda* Smith, el gusano cogollero, para el cual se utilizan anualmente 3,000 toneladas de ingrediente activo insecticida para su control; al cual le siguen otros lepidópteros como *Agrotis ipsilon* (Hufnagel), gusano trozador, y *Helicoverpa zea* Boddie, gusano elotero, los que en su conjunto se les aplica insecticida 1–3 veces en cada ciclo de cultivo. Científicos mexicanos han trabajado en diferentes aspectos del MIP en maíz; sin embargo estas tácticas casi nunca son utilizadas por los agricultores mexicanos. Un grupo de científicos y asesores agrícolas en México fue entrevistado para obtener la importancia de las plagas del maíz identificando la presencia temporal y territorial de éstas durante las diferentes etapas de desarrollo del cultivo. Se obtuvo también un inventario de las prácticas de control más usadas, las dosis y frecuencia con las que se hacen las aplicaciones de insecticida. Esta información, no disponible anteriormente, intenta apoyar a investigadores, funcionarios de gobierno y especialistas de la industria a priorizar esfuerzos y recursos para el diseño e implementación de técnicas, prácticas y programas MIP para la producción de maíz en México.

**Key Words:** *Spodoptera*, maize production, insecticide use, temporal and geographical pest occurrence

Maize (*Zea mays* L.) is the most widely produced crop in the world (U.S. Department of Agriculture, Foreign Agricultural Service [USDA-FAS] 2013). Mexico, the fifth largest maize producer and area of origin and domestication of this crop (Matsuoka et al. 2002), has planted annually ≈6.8 million hectares in the past 5 yr, representing 30% of its total domestic agricultural production and 6.6% of the country's arable land (Secretariat of agriculture, livestock, rural development, fisheries and food [SAGARPA] 2013, World Bank 2013). However, the country's average yield of 3.17 tons per hectare is 38% below the world's average (USDA-FAS 2013).

Many factors limit maize yield and productivity around the world. Weeds, the main pest problem, reduce yield potential by as much as 11%; animal pests contribute with 10% of the reduction and plant pathogens an additional 11% (Oerke 2006). Although maize is a major crop globally, it is typically not grown under modern or comprehensive

integrated pest management (IPM) programs. IPM is a strategy to avoid or minimize economic damage and to lower environmental impact of pest control relative to the indiscriminate use of pesticides. Multiple tools and tactics based on the biology and population ecology of crops and pests, economic feasibility of the crop, and decision and action thresholds, are incorporated into an IPM program (Higley and Pedigo 1996, Kogan 1998, U. S. Environmental Protection Agency [U.S. EPA] 2013). As a reference, in 1986, only 20% of the maize crop area in the United States, the largest producer in the world (U.S. Department of Agriculture, National Agricultural Statistics Service [USDA-NASS] 2013), had adopted this pest control management (Morse and Buhler 1997). The global situation has not changed significantly in recent years especially in Mexico. Approximately 3,000 tons of active ingredient per year are used to combat the fall armyworm (*Spodoptera frugiperda* Smith) (Blanco et al. 2010), the

most important maize pest in the American continent (Molina-Ochoa et al. 2001, Morillo and Notz 2001, Murúa and Virla 2004, de Melo et al. 2006, Zenner de Polanía et al. 2009). The severity of this pest in maize and lack of IPM program in many important crops has placed Mexico as the country using the highest quantity of pesticides per arable land in North America (Stokstad 2013).

Based on their level of technology adoption, maize production systems in Mexico may be classified as “commercial” maize, usually involving high capital investment and more technology, and “self-consumption” maize, involving high manual labor investment and less technology (SAGARPA 2013). Damaging pest populations occur in all production systems across the country but are mainly managed and controlled in commercial maize, not as much in self-consumption maize. Arthropod pest management in commercial maize relies primarily on chemical control using conventional synthetic insecticides.

Mexican farmers have successfully adopted IPM programs for tomatoes (Trumble and Alvarado-Rodríguez 1993), pecan trees (Rodríguez del Bosque and Tarango-Rivero 1997), broccoli and chili peppers (Blanco 2005), and for fruit flies (Tephritidae) (Mota-Sanchez et al. 2003). Different IPM components including biological control agents have been identified for all major maize pests. These include nearly 40 diseases, predators, and parasitoids of fall armyworm (Molina-Ochoa et al. 2001, Cortez-Mondaca et al. 2012, Estrada-Virgen et al. 2013), nine parasitoids of the leafhopper *Dalbulus* spp. (Moya-Raygoza et al. 2008), 13 parasitoids of the bollworm–tobacco budworm (*Helicoverpa zea*–*Heliothis virescens*) complex (Loera-Gallardo et al. 2008), and 38 of *Diatraea* spp., and *Eoreuma loftini* (Dyar) borers (Rodríguez del Bosque and Vejar-Cota 2008). Natural enemies are able to keep only 10% or less of *S. frugiperda* populations below the common action threshold of  $\geq 20\%$  of the plants with *S. frugiperda* larvae (Cruz and Turpin 1983, Harrison 1984, Fernández 2002). Control alternatives, such as the use of beneficial nematodes (Molina-Ochoa et al. 1999), bio-rational and low-impact insecticides (Méndez et al. 2002, García-Gutiérrez et al. 2012), and different formulations and dosage of selective insecticides (Castillejos et al. 2002, Zamora et al. 2008, Rosas-García et al. 2009) are examples of some of the published research. Although “bio-rational” insecticides have been shown effective and are available in the domestic market, their current use in Mexico is negligible, in part due to relative cost. For example, retail price of a *Bacillus thuringiensis*-based insecticide is often twice as much as that of a synthetic insecticide; these products are only used in  $\approx 1.5\%$  of the maize crop area grown in Mexico (Tamez Guerra et al. 2001). This type of relevant IPM information, however, is mostly published in academic journals where it is not accessible to most growers.

Perhaps the greatest obstacle for full implementation of maize IPM programs in Mexico is the large number and great diversity of growers. Approximately 2 million growers plant maize under very diverse conditions across Mexico; 28% of the 1.8 million maize fields in Mexico are between 1 and 2 ha in area (SAGARPA 2013). This makes it a great challenge to deliver practical information and appropriate solutions. An IPM program represents a holistic approach that requires recognizing and quantifying biological control agents and pests, to have knowledge of pesticides, and to adopt better adapted plant varieties for specific regions. What proves even more difficult is coordinating all these concepts and actions with crop irrigation and tillage, planting schedules, fertilization and harvesting, and to have knowledge of the markets, among other practices (Morse and Buhler 1997). Solid IPM programs require continuing education because pest species, population dynamics, and action threshold, considered the “pillars” of IPM (U.S. EPA 2013), can vary significantly as new knowledge and technologies become available, as new pests emerge, and as climate and economic outlook changes. Morse and Buhler (1997) point out that perhaps an IPM program corresponds more to the characteristics of the researchers and academics that develop these programs, than to the actual needs of the grower, as the former tend

to specialize in a given discipline, or even worse, in a specific part of the problem, leaving the remaining parts of the development and implementation of an IPM program to other colleagues (Morse and Buhler 1997). For growers to evaluate the impact of each of the IPM components and adopt them, they require highly specialized knowledge of diverse disciplines. This may be overwhelming and can lead to adoption of more user-friendly and quicker solution to the problem, which often implies the use of pesticides, without taking into account all the factors that may justify this practice. Growers can be persuaded initially to adopt an IPM program through a financial analysis and later on, regulatory and environmental considerations may dictate the extent to which programs are adopted (Koul and Cuperus 2007). A critical financial analysis is key for an IPM program, and the lack of interdisciplinary considerations in these analyses is generally not discussed (Lipke et al. 1987). It is not surprising that most of the information on IPM flows through the agrochemical suppliers and distributors (Koul and Cuperus 2007), which may explain why pest control consists mostly of pesticide use.

Most of the information on maize pests and IPM in Mexico has been published in technical bulletins of a reduced regional scope or in difficult-to-access reports, generally describing laboratory screenings, preliminary field evaluations, or listing technical recommendations, but rarely include products and practices that growers actually use. The goal of this report is to provide up-to-date information on the common pests in main maize-growing areas of Mexico from experts in the field. Updated information on maize pest population dynamics and control has been gathered from agricultural professionals directly involved in maize crop protection. This critical information, not readily available for Mexican production systems, is intended to help researchers, government officials, and industry specialists in making decisions regarding time and resource allocations, and in the design and implementation of IPM techniques, practices, and programs in Mexican maize.

## Approach and Methods

Between 2010 and 2013 an expert panel composed of Mexican researchers and crop advisors gathered information regarding the main maize pests that reduce maize production and require control. Expert panels have been used successfully to gather base line data in the absence of formal documentation (Department of Environment and Primary Industries [DEPI] 2014). The geographical areas where information was obtained included commercial and self-consumption production systems and maize grain and silage production, information based on statistics obtained from the Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación (Secretariat of agriculture, livestock, rural development, fisheries and food; SAGARPA 2013; Table 1).

Detailed information was gathered from agricultural specialists who provided data from commercial maize in different regions of Mexico. Responses to a common questionnaire were obtained from 24 participants including established researchers and technical personnel who were asked to—1) list the important arthropod pests that cause yield loss in the region; 2) identify the developmental stages of maize in which each pest species causes crop injury; 3) list the types of control strategies used for each pest or pest complex and its relative cost; and 4) discuss the kind of integrated crop and pest management tactics used (if any) and their elements.

Answers were received from experts representing seven growing regions: 1) “Bajío,” which includes the states of Guanajuato, Michoacán, and Querétaro; 2) Chiapas state; 3) Chihuahua, where data were obtained from five different regions within the state; 4) “Laguna,” including data from Coahuila and Durango states; 5) “Pacific,” with information from Colima, Jalisco, and Nayarit states; 6) “North-west,” the southern portion of Sonora and the northern part of Sinaloa states; and 7) Tamaulipas state, north and south (Table 1). These seven regions represent almost half (45%) of the maize grain production area

**Table 1. Corn production in Mexico in 2011 in selected regions and nation-wide statistics**

	Spring–summer planting				Autumn–winter planting			
	Irrigated		Rain fed		Irrigated		Rain fed	
	Hectares	Yield <sup>a</sup>	Hectares	Yield <sup>a</sup>	Hectares	Yield <sup>a</sup>	Hectares	Yield <sup>a</sup>
	Grain production							
Bajío	220,648	7.12	446,137	1.47	10,421	4.37	142	2
Chiapas	1,200	3.19	590,096	2.28	10,120	4.47	105,027	1.5
Chihuahua	97,925	8.6	20,546	0.45	0		0	
Laguna	30,205	3.98	64,361	0.37	49	3.5	0	
Northwest	10,913	4.77	32,217	0.9	392,518	6.66	3,622	2.5
Pacific	38,316	5.94	523,415	4.04	8,620	5.35	3,808	4
Tamaulipas	6,462	2.2	26,624	1.13	89,726	4.94	1,616	1.32
COUNTRY	668,183	5.91	4,392,199	2.11	578,677	6.41	430,033	1.67
	Silage production							
Bajío	16,506	55.92	50	11	134	22	0	
Chiapas	0		0		0		0	
Chihuahua	13,250	38.82	10,857	5.21	0		0	
Laguna	39,265	47.03	24,895	5.98	0		0	
Northwest	629	14.08	320	8.06	0		0	
Pacific	19,902	27.59	80,523	29.42	632	27.85	943	19.82
Tamaulipas	0		0		0		0	
COUNTRY	130,646	49.01	177,794	17.55	1356	32.16	1,430	26.06

Information source: [http://siap.gob.mx/index.php?option=com\\_wrapper&view=wrapper&Itemid=351](http://siap.gob.mx/index.php?option=com_wrapper&view=wrapper&Itemid=351)  
<sup>a</sup> Thousands of kilograms per hectare.

**Table 2. Presence of the fall armyworm, *S. frugiperda*, larvae in corn in different corn-growing regions of Mexico indicating the crop development stage**

<sup>a</sup>	Dec.	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sep.
1 Bajío						V2	V4	V6		
1 Chiapas								V2	V12	V2
1 Chih SE					V2	V4	V6	V8		
1 Chih NE						V2	V4	V6	V8	
1 Chih N							V2	V4	V6	V8
2 Laguna				V2	V4	V6	V8		V2	V4
1 Northwest	V2	V3	V4		V2	V4				
2 Pacific							V2	V4	V6	
1 Tamp N				V2	V4	V6	V8	V12		
1 Tamp S		V2	V4	V6	V12	R1			V2	V4
									V10	VT
										R2


Corn development stages according to <http://weedsoft.unl.edu/documents/growthstagesmodule/corn/corn.htm>.  
<sup>a</sup> Pest problem ranking.

(2,734,734 hectares) and two thirds (67%) of the maize production in the country (Table 1).

Additional information on the use of insecticides on maize and growers' preferred products was obtained from a confidential pesticide market research study conducted in 2009, commissioned by an agrochemical company, which was used only as comparison for our data. Calculation of the environmental impact of pesticides (Kovach et al. 1992) was carried out through the Cornell University's web page (Cornell University 2014). Information of peer-reviewed papers on the subject published in accredited journals from 2000 to 2012 was obtained on the World Wide Web (Internet) through the Biosis search engine using the key words "pests" and "Mexico," including only those reports where the authors were affiliated with a Mexican institution.

**Results and Discussion**

*S. frugiperda* is the main pest of maize in five of the seven regions, nearly 3 million hectares in our survey region, representing 45% of all maize planted in Mexico (Table 1). In the other two regions (Laguna and Pacific), *S. frugiperda* was the second most important pest infesting all 2.9 million hectares represented in this survey (Table 2; Fig. 1). The fall armyworm is controlled with two to three applications of insecticide (Table 3). According to a survey carried out in Mexico by a marketing company in 2009 (Dow AgroSciences, personal communication), including maize farmers in all the regions of the country,



**Fig. 1.** Hectares affected by different maize pests in Mexico. FAW, *Spodoptera frugiperda*; CW, *Agrotis ipsilon*; WG, *Phyllophaga* spp., and other white grubs; THR, *Frankliniella* sp./*Thrips tabaci*; TSM, *Tetranichus urticae*; CEW, *Helicoverpa zea*; GA, *Sitobion avenae*; CRW, *Diabrotica* spp.; CLA, *Rhopalosiphum maidis*; FB, *Epitrix* sp.; BAW, *Spodoptera exigua*.

growers agreed that the fall armyworm is the most difficult pest to control.

There are no good examples of comprehensive IPM programs for maize in any of the regions studied in Mexico. Although Mexican

Table 3. Insecticides used to control different corn pest in Mexico

	Rate (g AI/ha)	EIQ <sup>a</sup>	Bajío (2)	Chiapas (2)	Chih SE (2)	Chih NE (2.5)	Chih N (2)	Laguna (2)	Pacific (2)	Northwest (2)	Tamp N (2)	Tamp S (2)
Abamectin	11.1	0.3			BGM (2.5) TSM (1.5)		TSM (1)	BGM (2.5) TSM (1.5)		TSM (1)		
Acephate	1,088	24.9					FAW (2), CEW (1.5)					
<i>B. thuringiensis</i> <sup>b</sup>	52 <sup>b</sup>	7.2					BAW (<0.5)					
Carbofuran	1,000	45.2	WG (1–1.5)		CW (1)		WG (<0.5)		FAW (2)			
Chlorfenapyr	120	1.6					CEW (1.5)		WG (2)			
Chlorpyrifos	480	11	FAW (2), WG (1.5), Nre (2), LMB (2)	FAW (2), CW (1)	FAW (2), CW (1)	CEW (2.5), BAW (1)	FAW (2), CEW (1)	FAW (2), CW (1)	FAW (2), WG (2), BAW (1)	FAW (2), CEW (1), CW (1)		FAW (2)
Diazinon	1,000	39.3					WG (<0.5)					
Diflubenzuron	48	1.0					BAW (<0.5)					
Dicofol	368–613	9.4					TSM (0.5–1)			TSM (1)		
Dimethoate	435	12.5	THR (1)					FB (1.5), CRW (1.5), THR (1), CLA (1)				
Emamectin benzoate	10–16	2.2	FAW (2)				FAW (2), CEW (1.5)			FAW (2), CEW (1)		FAW (2)
Fenpyroximate	75	1.2										
Methamidophos	841	26.5					FAW (2), CEW (1)			GA (1)		
Methomyl	450–508	8.5						BGM (2.5) TSM (1.5) FB (1.5), CRW (1.5), THR (1)		FAW (2), CEW (1), CW (1)		
Methyl parathion	720	13.1	Nre (2), LMB (2)									
Novaluron	44–88	0.5							FAW (2)			
Omethoate	400	11.5							WG (2)			
Phorate	1,125	49										
Propargite	1,178–2,731	106										
Pyrethroids <sup>b</sup>	35–188	1.1	FAW (2), Nre (2), LMB (2)	FAW (2), CW (1)	FAW (2)	CEW (2.5), BAW (1)	TSM (0.5)	FAW (2)	FAW (2), BAW (1)	FAW (2), CW (1)	FAW (2), CEW (1)	FAW (2), CEW (1)
Rynaxypyr	50	NA										
Spinetoram	Jun-50	1.2	FAW (2)									FAW (2)
Spinosad	35	0.4										FAW (2)
Spiromesifen	120	2.9										
Terbufos	1,000	58.9	WG (1–1.5)									
Thiodicarb	235	4.7										

The number in parenthesis refer to the average number of applications per pest in a particular region.

BAW, *S. exigua*; CEW, *H. zea*; CRW, *D. virgifera*; CW, *A. ipsilon*; FAW, *S. frugiperda*; FB, flea beetle (*Epitrix* sp.); GA, *Sitobion avenae*; LMB, *Nicentrites testaceipes*; THR, Thrips; BGM, *Oligonychus pratensis*; TSM, *T. urticae*; WG, *Phyllophaga* and other white grubs; CLA, *Rhopalosiphum maidis* and *Sitobion avenae*.

<sup>a</sup> Environmental impact (EIQ) calculated for one application at its lowest recommended rate from: <http://ceeeiq-lamp.cit.cornell.edu/EIQcalc/input.php?cat>.

<sup>b</sup> Calculated for Dipel 54% and expressed as *B. thuringiensis* fermentation solids, spores, and insecticidal toxins.

<sup>c</sup> Calculated for alpha cypermethrin, the most commonly used pyrethroid.



**Table 5. Presence of coleopteran pests in corn in different corn-growing regions of Mexico indicating the crop developmental stage**

<sup>a</sup>	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	
<i>Phyllophaga</i> spp. and other white grubs												
2 Bajío					VE	V2	V4					
9 Chih SE						V2	V2	V4				
6 Chih N						V2	V2	V4				
1 Pacific					S	VE	V2	V4				
Corn rootworms, <i>Diabrotica</i> spp.												
6 Chih SE						V8	V12	VT	R1	V8	V12	
5 Chih N							V8	V12	VT	R1		
6 Laguna					V8	V12	VT	R1	V8	V12	VT	R1
Flea beetles, <i>Epitrix</i> sp./ <i>Chaetocnema</i> sp.												
3 Chih SE						V8	V12	VT	R1			
3 Laguna					V8	V12	VT	R1	V8	V12	VT	R1
Lesser maize billbug, <i>Nicentrites testaceipes</i> (Champion)												
3 Bajío							VT	R1				

Corn development stages according to <http://weedsoft.unl.edu/documents/growthstagesmodule/corn/corn.htm>.

<sup>a</sup> Pest problem ranking.

networks that may provide information on alternatives to broad-spectrum pesticides, which can greatly influence decision making in pest management (Koul and Cuperus 2007).

In addition to fall armyworm, other important pests of Mexican maize include mites *Oligonychus pratensis* (Banks) and *Tetranychus urticae* Koch primarily damaging silage maize in Laguna and parts of Chihuahua (Table 6) and affecting almost 1.5 million hectares of the area of our survey (Fig. 1), and the *Phyllophaga* spp. white grub complex in the Pacific and Bajío regions (Table 5), with similar geographic impact (Fig. 1). Other Lepidoptera pests are also very important in Mexico. The black cutworm, *Agrotis ipsilon* (Hufnagel), is controlled with insecticides on 2.2 million hectares (Fig. 1). However, this insect is relatively easy to manage (one insecticide application, Table 3); therefore, respondents to our survey ranked it as second to fifth in importance (Table 4). *Helicoverpa zea* (Boddie) is considered a serious problem in four of the seven regions (Table 7), impacting 700,000 hectares (23%), requiring between one and three sprays per crop cycle (Table 3). The beet armyworm, *Spodoptera exigua* (Hübner), and the Mexican rice borer, *E. loftini*, were considered among the five most harmful pests (Table 4). Although *S. exigua* only affects 100,000 ha of our survey area (Fig. 1), it requires one to three insecticide applications (Table 3).

Host plant resistance has been used to reduce the impact of Lepidoptera pests, mainly *S. frugiperda* in Mexico. Zapalote Chico and Antigua varieties have shown to be resistant to fall armyworm, corn earworm, and other Lepidoptera pests that attack different parts of the plant (Williams et al. 1978, Wiseman and Davis 1990, Abel et al. 2000). Maize host plant resistance efforts often rely on recombinant DNA techniques to develop genetically modified (GM) hybrids (mainly expressing proteins of *Bacillus thuringiensis*) and ribonucleic acid interference (RNAi), effective strategies against Lepidoptera and Coleoptera pests (Lynch et al. 1999, Nuessly et al. 2007, Price and Gatehouse 2008). Pest-resistant maize derived from crosses with Zapalote Chico or Antigua is not very common in commercial hybrids currently available, likely due to emphasis on development of GM varieties (W. P. Williams, 2013, personal communication). A clear advantage of recombinant DNA technologies that produce GM varieties over traditional ones was determined by comparing the performance of a GM *B. thuringiensis*-expressing variety with a conventional maize variety to which pyrethroids were applied eight times (>200 g ai/ha) to protect from a heavy infestations of *S. frugiperda* and *H. zea*. The GM hybrid with no synthetic insecticide applications produced 100% ears free from insect damage, whereas the conventional hybrid treated multiple times with insecticide produced only

**Table 6. Presence of several pests in corn in different corn-growing regions of Mexico indicating the crop developmental stage**

<sup>a</sup>	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.		
<i>Banks</i> grass mite, <i>Oligonychus pratensis</i> /two-spotted spider mite, <i>Tetranychus urticae</i>												
2 Chih SE						V8	V12	VT	R1			
4 Chih NE						V8	V12	VT	R1			
3 Chih N							V8	V10	V12	VT	R1	
1 <sup>b</sup> Laguna					V8	V12	VT	R1	V8 <sup>b</sup>	V12	VT	R1 <sup>b</sup>
6 Northwest			V4	R1								
Corn leaf aphid, <i>Rhopalosiphum maidis</i> Fitch												
7 Chih SE						V8	V12	VT	R1			
5 Chih NE						V8	V12	VT	R1			
7 Laguna					V8	V12	VT	R1	V8	V12	VT	R1
Grain aphid, <i>Sitobion avenae</i>												
4 Chiapas									R1	R1		
4 Northwest	R1	R1		R1	R1							
Thrips, <i>Frankliniella</i> sp./ <i>Thrips tabaci</i>												
4 Bajío					VE	V2						
4 Chih SE						V8	V12	VT	R1			
4 Laguna					V8	V12	VT	R1	V8	V12	VT	R1
4 Northwest			VE	V2								
Corn leafhopper, <i>Dalbulus maidis</i> (De Long & Wolcott)												
5 Chih N								VT	R1			

Corn development stages according to <http://weedsoft.unl.edu/documents/growthstagesmodule/corn/corn.htm>.

<sup>a</sup> Pest problem ranking.

<sup>b</sup> In the summer corn planting this pest is ranked number 8.

**Table 7. Presence of the corn earworm, *Helicoverpa zea*, larvae in corn in different corn-growing regions of Mexico indicating the crop developmental stage**

<sup>a</sup>		Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.
3	Chiapas							V8 V12	V8 V12		
8	Chih SE							R1			
2	Chih NE							VT R1			
2	Chih N								VT	R1	
7	Laguna						R1				R1
3	Pacific	R1	R1	V2 V4							
2	Tamp N				V8	V12					
3	Tamp S		V12	R1						R1 R2	

Corn development stages according to <http://weedsoft.unl.edu/documents/growthstagesmodule/corn/corn.htm>.  
<sup>a</sup> Pest problem ranking.

18% of ears with no fall armyworms and corn earworms (Shelton 2012). In another study comparing GM and conventional maize varieties, a 71% yield loss due to *S. frugiperda* attack was documented only in the conventional variety (Buntin et al. 2001). This level of performance has led to broad adoption of GM maize hybrids in different countries.

Data generated in Mexico have yielded similar results. GM maize compared with insecticide-treated (two to five applications ranging between 720–3,600 g ai/ha) conventional maize, showed 10% higher yields and effective control of fall armyworm and corn earworm infestations (Piña and Solleiro 2013). Reducing injury by *S. frugiperda* and *H. zea* to maize ears may also lead to reductions in the levels of aflatoxin (known carcinogen) produced by the fungus *Aspergillus flavus* (Gourama and Bullerman 1995, Boermans and Leung 2007, Williams et al. 2010).

However, the availability of improved hybrids developed through conventional or recombinant DNA technologies for small growers in Mexico faces great obstacles. Most of the small growers likely cannot afford the extra cost of insect-resistant hybrid seeds. Hybrid seed prices (≈US\$250.00/ha) represents almost half of the expected income from grain sales that a typical Mexican producer may obtain (not factoring government subsidies). The promising side of this situation is that seed companies have realized that even small growers are interested in improved varieties and in some areas of Mexico, maize hybrids are offered in small quantities (60,000 seeds per sack) to plant as little as one hectare. This indicates there is a viable market for improved maize seed for the small growers in Mexico. Interestingly, two or three insecticide applications per crop (Table 3) at a cost of US\$30–45/ha may be sufficient to keep the pest complex in check (not only Lepidoptera pests) in most instances. This is far below the cost of hybrid seed and may be particularly advantageous to the small farmer facing a diverse pest complex. Thrips, aphids, and mites were reported as important pests in almost half of the regions represented in our survey (Table 6) and cannot be effectively controlled by GM maize hybrids at this time. The future development of elite hybrids providing some type of cross-resistance and a significant increase in maize yields would drastically change the above equation.

Wide implementation of IPM of insects in Mexican maize is not a lost cause, but further delays in finding solutions to the low productivity challenge, which is partly a consequence of inadequate pest management, may force the country to continue importing ≈30% of its maize grain to keep up with internal demand (SAGARPA 2013). Lack of adoption of IPM tactics will likely lead Mexican maize growers to continue applying conventional pesticides to combat major pests (≈3,000 tons of active ingredient per year for a single pest; Blanco et al. 2010).

Possible solutions to these problems may be achieved by “1) a change in pest management approaches through demonstration-research plots where growers can verify the benefits of IPM programs,” along with public campaigns. Similar programs have been

successful for other crops in Asia where marked reductions in insecticide use and higher crop yields have been documented (Normile 2013); “2) an easy and accessible way for growers to recognize and understand the function of natural enemies and the benefits of bio-rational alternatives in minimizing negative impact of pesticides on beneficial organisms.” As the use of the Internet becomes more common, information dissemination becomes easier. There are different sites available on the World Wide Web containing relevant information, but a better portal should be created to link the work of Mexican professionals directed at solving specific pest problems; “3) development and marketing of pest-resistant maize hybrids through effective and affordable methods to reduce the need for broad-spectrum pesticides.” This should include a science-based assessment on the impact of agricultural biotechnology taking advantage of the great maize genetic diversity present in Mexico to identify appropriate sources of host plant resistance; “4) development and marketing of bio-rational products and biological control products for the small farmer” (smaller, lower cost packaging). This should also involve greater knowledge dissemination of the ecology of crops and herbivores for better managing the latter; and “5) other elements” including training of involved technical personnel on technology transfer and spreading of information, on government policies reflecting present day production of maize and on assessment and ongoing evaluation of programs (Council for Agricultural Science and Technology [CAST] 2003).

#### Acknowledgments

We thank W. Paul Williams for his advice on some aspects of this report, and Drs. Susan T. Ratcliffe, Craig A. Abel, Michael Wach, Julio S. Bernal, Carlos Bográn, Fred Gould, Susana G. Fredín, and an anonymous reviewer for their editorial comments.

#### References Cited

- Abel, C. A., R. L. Wilson, B. R. Wiseman, W. H. White, and F. M. Davis. 2000. Conventional resistance of experimental maize lines to corn earworm (Lepidoptera: Noctuidae), fall armyworm (Lepidoptera: Noctuidae), southwestern corn borer (Lepidoptera: Crambidae), and sugarcane borer (Lepidoptera: Crambidae). *J. Econ. Entomol.* 93: 982–988.
- Armenta, R., A. M. Martínez, J. W. Chapman, R. Magallanes, D. Goulson, P. Caballero, R. D. Cave, J. Cisneros, J. Valle, V. Castillejos, et al. 2003. Impact of nucleopolyhedrovirus bioinsecticide and selected synthetic insecticides on the abundance of insect natural enemies on maize in southern Mexico. *J. Econ. Entomol.* 96: 649–661.
- Blanco, C. A. 2005. *A la hora de comer, ¿qué nos preocupa?*, vol. Fondo de Cultura Económica. Mexico D.F., Mexico.
- Blanco, C. A., M. Portilla, J. L. Jurat-Fuentes, J. F. Sánchez, D. Viteri, P. Vega-Aquino, A. P. Terán-Vargas, A. Azuara-Domínguez, J. D. López, R. Arias, et al. 2010. Susceptibility of isofamilies of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) to CryIAc and CryIFa proteins of *Bacillus thuringiensis*. *Southwest. Entomol.* 35: 409–415.
- Boermans, H. J., and M.C.K. Leung. 2007. Mycotoxins and the pet food industry: toxicological evidence and risk assessment. *Int. J. Food Microbiol.* 119: 95–102.

- Buntin, G. D., R. D. Lee, D. M. Wilson, and R. M. McPherson. 2001. Evaluation of YieldGard transgenic resistance for control of fall armyworm (Lepidoptera: Noctuidae) on corn. *Fla. Entomol.* 84: 37–42.
- (CAST) Council for Agricultural Science and Technology. 2003. Integrated pest management: current and future strategies. Council for Agricultural Science and Technology, Aimes, IA.
- Castillejos, V., J. Trujillo, L. D. Ortega, J. A. Santizo, J. Cisneros, D. I. Penagos, J. Valle, and T. Williams. 2002. Granular phagostimulant nucleopolyhedrovirus formulations for control of *Spodoptera frugiperda* in maize. *Biol. Control* 24: 300–310.
- Cornell University. 2014. A method to measure the environmental impact of pesticides. Cornell University, Ithaca, NY.
- Cortez-Mondaca, E., J. Pérez-Márquez, and F. Bahena-Juárez. 2012. Control biológico natural del gusano cogollero (Lepidoptera: Noctuidae) en maíz y sorgo, en el norte de Sinaloa, México. *Southwest. Entomol.* 37: 423–428.
- Cruz, I., and F. T. Turpin. 1983. Yield impact of larval infestations of the fall armyworm (Lepidoptera: Noctuidae) to midwhorl growth stages of corn. *J. Econ. Entomol.* 76: 1052–1054.
- de Melo, E. P., M. G. Fernandez, P. E. Degrande, R.M.A. Cessa, J. L. Salomão, and R. F. Nogueira. 2006. Distribuição espacial de plantas infestadas por *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) na cultura do milho. *Neotrop. Entomol.* 35: 689–697.
- (DEPI) Department of Environment and Primary Industries. 2014. Department of Environment and Primary Industries. Effective engagement: expert panel. <http://www.dse.vic.gov.au/effective-engagement/toolkit/tool-expert-panel>. Accessed in December 2013.
- Estrada-Virgen, O., J. Cambero-Campos, A. Robles-Bermudez, C. Ríos-Velasco, C. Carvajal-Cazola, N. Isordia-Aquino, and E. Ruíz-Cancino. 2013. Parasitoides y entomopatógenos nativos asociados al gusano cogollero *Spodoptera frugiperda* (Lepidoptera: Noctuidae) en Nayarit, México. *Southwest. Entomol.* 38: 339–344.
- Fernández, J. L. 2002. Nota corta: Estimación de umbrales económicos para *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) en el cultivo del maíz. *Invest. Agric. Prod. Prot. Veg.* 17: 467–474.
- García-Gutiérrez, C., M. B. González-Maldonado, and E. Cortez-Mondaca. 2012. Use of natural enemies and biorational pest control of corn. *Ra Ximhai* 8: 57–70.
- Gourama, H., and L. B. Bullerman. 1995. *Aspergillus flavus* and *Aspergillus parasiticus*: aflatoxigenic fungi of concern in foods and feeds: a review. *J. Food Prot.* 58: 1395–1404.
- Harrison, F. P. 1984. The development of an economic injury level for low populations of fall armyworm (Lepidoptera: Noctuidae) in grain corn. *Fla. Entomol.* 67: 335–339.
- Higley, L. G., and L. P. Pedigo. 1996. Economic thresholds for integrated pest management. University of Nebraska, Lincoln, NE.
- Kogan, M. 1998. Integrated pest management: historical perspectives and contemporary developments. *Annu. Rev. Entomol.* 43: 243–270.
- Koul, O., and G. W. Cuperus. 2007. Ecologically based integrated pest management: present concept and new solutions. CABI, Oxfordshire, United Kingdom.
- Kovach, J., C. Petzoldt, J. Degnil, and J. Tette. 1992. A method to measure the environmental impact of pesticides. *N.Y. Food Life Sci. Bull.* 139: 1–8.
- Lipke, L. A., H. W. Ladewig, and E. Taylor-Powell. 1987. National assessment of extension efforts to increase farm profitability through integrated programs. Texas Agricultural Extension Service, U.S. Department of Agriculture, TX.
- Loera-Gallardo, J., J. I. López-Arroyo, and M. A. Reyes-Rosas. 2008. Complejo *Heliothis virescens* y *Helicoverpa zea* (Lepidoptera: Noctuidae) pp. 57–74. Casos de control biológico en México. Multi Prensa, Mexico D. F., Mexico.
- Lynch, R. E., B. R. Wiseman, D. Plaisted, and D. Warnick. 1999. Evaluation of transgenic sweet corn hybrids expressing CryIA(b) toxin for resistance to corn earworm and fall armyworm (Lepidoptera: Noctuidae). *J. Econ. Entomol.* 92: 246–252.
- Matsuoka, Y., Y. Vigouroux, M. M. Goodman, J.S.G., E. Buckler, and J. Doebley. 2002. A single domestication for maize shown by multilocus microsatellite genotyping. *Proc. Natl. Acad. Sci. U.S.A.* 99: 6080–6084.
- Méndez, W. A., J. Valle, J. E. Ibarra, J. Cisneros, D. I. Penagos, and T. Williams. 2002. Spinosad and nucleopolyhedrovirus mixtures for control of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in maize. *Biol. Control* 25: 195–206.
- Molina-Ochoa, J., R. Lezama-Gutiérrez, J. J. Hamm, B. R. Wiseman, and M. Lopez-Edwards. 1999. Integrated control of fall armyworm (Lepidoptera: Noctuidae) using resistant plants and entomopathogenic nematodes (Rhabditida: Steinernematidae). *Fla. Entomol.* 82: 263–271.
- Molina-Ochoa, J., J. J. Hamm, R. Lezama-Gutiérrez, M. Lopez-Edwards, and M. Gonzalez Ramirez. 2001. A survey of fall armyworm (Lepidoptera: Noctuidae) parasitoids in the Mexican states of Michoacán, Colima, Jalisco, and Tamaulipas. *Fla. Entomol.* 84: 31–36.
- Molina-Ochoa, J., J.E.J. Carpenter, R. Lezama-Gutiérrez, J. E. Foster, M. González-Ramírez, C. A. Angel-Sahagún, and J. Fariás-Larios. 2004. Natural distribution of hymenopteran parasitoids of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) larvae in Mexico. *Fla. Entomol.* 87: 461–472.
- Morillo, F., and A. Notz. 2001. Resistencia de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) a lambdacialotrina y metomil. *Entomotropica* 16: 79–87.
- Morse, S., and W. Buhler. 1997. Integrated pest management: ideals and realities in developing countries. Lynne Rienner Publishers, Boulder, London, United Kingdom.
- Mota-Sanchez, D., F. Santos Gonzalez, B. Alvarado-Rodriguez, O. Diaz-Gomez, H. Bravo-Mojica, G. Santiago-Martinez, and R. Bujanos. 2003. Integrated Pest Management in Mexico, pp. 273–284. Integrated Pest Management in the Global Arena. CAB International Publishing, United Kingdom.
- Moya-Raygoza, G., N. M. Chacón-Torres, V. Palomera-Avalos, I. M. Becerra-Chiron, and J. S. Bernal. 2008. Effects of Corn Stunt Spiroplasma on Survival of *Dalbulus maidis* (Hemiptera: Cicadellidae) Parasitized or Preyed upon by *Gonatopus bartletti* (Hymenoptera: Dryinidae) or Infected by *Metarhizium anisopliae*. *Ann. Entomol. Soc. Am.* 101: 113–118.
- Murúa, M. G., and E. G. Virla. 2004. Presencia invernal de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) en el área maicera de la provincial de Tucumán, Argentina. *Rev. de la Facultad de Agronomía, La Plata* 105: 46–52.
- Normile, D. 2013. Vietnam turns back 'a tsunami of pesticides'. *Science* 341: 737–738.
- Nuessly, G. S., B. T. Scully, M. G. Hentz, R. Beiriger, M. E. Snook, and N. W. Widstrom. 2007. Resistance to *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and *Euxesta stigmatias* (Diptera: Ulidiidae) in sweet corn derived from exogenous and endogenous genetic systems. *J. Econ. Entomol.* 100: 1887–1895.
- Oerke, E. C. 2006. Crop losses to pests. *J. Agric. Sci.* 144: 31–43.
- Piña, S., and J. L. Solleiro. 2013. México, pp. 341–410. Introducción al ambiente del maíz transgénico: análisis de ocho casos en Iberoamérica. D.R. AGROBIO México.
- Price, D.R.G., and J. A. Gatehouse. 2008. RNAi-mediated crop protection against insects. *Trends Biotechnol.* 26: 393–400.
- Rodríguez del Bosque, L., A., and S. H. Tarango-Rivero. 1997. Manejo integrado de plagas del nopal. Fundación Produce Chihuahua, Mexico.
- Rodríguez del Bosque, L. A., and G. Vejar-Cota. 2008. Barrenadores del tallo (Lepidoptera: Crambidae) del maíz y caña de azúcar, pp. 9–22. Casos de control biológico en México. Multi Prensa, Mexico, D. F., Mexico.
- Rosas-García, N. M., J. M. Villegas-Mendoza, and J. A. Torres-Ortega. 2009. Design of a *Bacillus thuringiensis*-based formulation that increases feeding preference on *Spodoptera exigua* (Lepidoptera: Noctuidae) larvae. *J. Econ. Entomol.* 102: 58–63.
- (SAGARPA) Secretariat of agriculture, livestock, rural development, fisheries and food. 2013. Producción agropecuaria. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación.
- Shelton, A. M. 2012. Genetically engineered vegetables expressing proteins from *Bacillus thuringiensis* for insect resistance: Successes, disappointments, challenges and ways to move forward. *GM Crops Food Biotechnol. Agric. Food Chain* 3: 1–9.
- Stokstad, E. 2013. Pesticide planet. *Science* 341: 730–731.
- Tamez Guerra, P., L. J. Galán Wong, H. Medrano Roldán, C. García Gutiérrez, C. Rodríguez Padilla, R. A. Gómez Flores, and R. S. Tamez Guerra. 2001. Bioinsecticidas: su empleo, producción y comercialización en México. *Ciencia U.A.N.L.* 4: 143–152.
- Thompson, L. 2013. Pesticide impacts on beneficial species. Grape and wine research and development corporation. <http://research.agwa.net.au/wp-content/uploads/2012/09/2012-05-FS-Pesticide-Impacts2.pdf>. Accessed in October 2013.
- Trumble, J. T., and B. Alvarado-Rodríguez. 1993. Development and economic evaluation of an IPM program for fresh market tomato production in Mexico. *Agric. Ecos. Environ.* 43: 267–284.
- (USDA-FAS) U.S. Department of Agriculture, Foreign Agricultural Service. 2013. World agricultural production, Circular Series WAP 3–13, March 2013. U.S. Department of Agriculture, Foreign Agricultural Service. <http://usda01.library.cornell.edu/usda/fas/worldagproduction/2010s/2013/worldag-production-04-10-2013.pdf>. Accessed in July 2013.
- (USDA-NASS) U.S. Department of Agriculture, National Agricultural Statistics Service. 2013. Statistics by subject. U.S. Department of Agriculture, National Agricultural Statistics Service. [http://www.nass.usda.gov/Statistics\\_by\\_Subject/](http://www.nass.usda.gov/Statistics_by_Subject/). Accessed in July 2013.

- (U.S. EPA) Environmental Protection Agency. 2013. Integrated Pest Management (IPM) Principles. U.S. Environmental Protection Agency. <http://www.epa.gov/opp00001/factsheets/ipm.htm>. Accessed in October 2013.
- Williams, W. P., F. M. Davis, and G. E. Scott. 1978. Resistance of corn to leaf-feeding damage by the fall armyworm. *Crop Sci.* 18: 861–863.
- Williams, W. P., G. L. Windham, M. D. Krakowsky, B. T. Scully, and X. Ni. 2010. Aflatoxin accumulations in Bt and non-Bt maize testcrosses. *J. Crop Improv.* 24: 392–399.
- Wiseman, B. R., and F. M. Davis. 1990. Plant resistance to insects attacking corn and grain sorghum. *Fla. Entomol.* 73: 446–458.
- World Bank. 2013. Agricultural land (% of land area). The World Bank. <http://data.worldbank.org/indicator/AG.LND.AGRI.ZS>. Accessed in July 2013.
- Zamora, M. C., A. M. Martínez, M. S. Nieto, M. I. Schneider, J. I. Figueroa, and S. Pineda. 2008. Acticity of several biorational insecticides against the fall armyworm. *Rev. Fitotec. Mex.* 31: 351–357.
- Zenner de Polanía, I., H. A. Arévalo-Maldonado, R. Mejía-Cruz, and J. L. Díaz-Sánchez. 2009. Spodoptera frugiperda: respuestas de distintas poblaciones a la toxina Cry1Ab. *Rev. Colomb. Entomol.* 35: 34–41.

*Received 19 March 2014; accepted 8 September 2014.*